



International Journal of Chemistry and Pharmaceutical Sciences

Journal Home Page: www.pharmaresearchlibrary.com/ijcps



Research Article

Open Access

QSAR Studies of Some Substituted Pyrazole Derivatives against *S.aureus*

Kishor Arora^{*1} and Veena Nathani²

¹Department of Chemistry, Govt. P.G. College (Auto.) Datia, M.P.) India

²Department of Chemistry, Govt. K.R.G. Auto. P.G. College, Gwalior, M.P, India

ABSTRACT

In the present paper QSAR studies related to some novel substituted pyrazole derivatives are reported. All the compounds were evaluated for antimicrobial activity against *S.aureus*. Their reported antimicrobial activities were used for Quantitative Structure Activity Relationship (QSAR) studies to find correlation between different calculated molecular descriptor of the compounds and biological activity.

Keywords: Antimicrobial activity, *S.aureus*, Pyrazoles, QSAR studies

ARTICLE INFO

CONTENTS

1. Introduction	2043
2. Experimental	2044
3. Results and Discussion	2044
4. Conclusion	2048
5. References	2048

Article History: Received 14 August 2015, Accepted 21 September 2015, Available Online 27 October 2015

*Corresponding Author

Kishor Arora
Department of Chemistry,
Govt. P.G. College (Auto.)
Datia, M.P, India
Manuscript ID: IJCPS2712



PAPER-QR CODE

Citation: Kishor Arora. QSAR Studies of Some Substituted Pyrazole Derivatives against *S.aureus*. *Int. J. Chem, Pharm, Sci.*, 2015, 3(10): 2043-2049.

Copyright© 2015 Kishor Arora. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original work is properly cited.

1. Introduction

One of the key objectives of organic and medicinal chemistry is to design and synthesize molecule that possess potent therapeutic values. The rapid development of resistance to existing antimicrobial drugs generates a serious challenge to the scientific community. Consequently, there is a vital need for the development of International Journal of Chemistry and Pharmaceutical Sciences

new antimicrobial agents with potent activity against resistant microorganism [1-15]. Pyrazole derivatives have a long history of application in agrochemicals as herbicides and insecticides and in pharmaceutical industry. Due to its wide range of biological activity, pyrazoles have received a considerable interest in the field of drug discovery and

therefore pyrazole ring constitute a relevant synthetic target in pharmaceutical industry. In fact, such heterocyclic compounds represent the core structure of a number of drugs.

QSAR, one of the methods used to correlate the biological property of molecule with molecular descriptor derived from chemical structure. It is a mathematical model of a statistically validated correlation between the chemical structure and their activity profile [16].

Prompted by the above facts and in continuation to our efforts in the study of novel compounds for antimicrobial infection, we hereby report the quantum chemical based QSAR studies of pyrazole compounds.

2. Experimental

Twelve pyrazole derivatives have been identified for their antibacterial activity against *S.aureus*. All the compounds were screened for antibacterial activity at 200µg/ml concentration. Their reported antibacterial activities are mentioned in table 1.

Table 1: Antibacterial activities

Compound	<i>S.aureus</i>
A-1	16
A-2	23
A-3	18
A-4	18
A-5	14
A-6	25
A-7	24
A-8	22
A-9	20
A-10	24
A-11	26
A-12	13

Computational details:

AM1 and PM3 Hamiltonia were studied for these compounds to develop one dimensional and later three dimensional descriptors. The structures of compounds were drawn using professional version of HYPERCHEM software 8.0. The descriptors studied were Surface Area (SAA), Surface Area Grid (SAG), Volume (VOL), Hydration Energy (HE), Refractivity (REF), Polarizability (POL), Total Energy (TE), Electronic Energy (EE), Heat of Formation (HF), Dipole Moment (DM) and Zero Point Energy (ZPE). The calculations were conducted with the aid of Pentium core-2 duo machine with the following configuration

Intel @ core TM 2 Duo CPU
T5450@1.66GHz
982 MHz 896 MB RAM
150 GB HDD

Windows - Microsoft windows XP software as an operating system. Regression analyses to get QSAR equations were performed and statistical calculations were done with the help of MS EXCEL software.

3. Results and Discussion

Analytical studies of compounds

The compounds taken for the studies were reported earlier and their structural viz. analytical spectral and other studies were also reported. Their structures were established and taken as such for the studies in the present paper. The analytical studies related to these compounds are reported in table 5.

Antibacterial activity

All the compounds were screened for antibacterial activity. However, compound A-2, A-3, A-6, A-10 and A-11 have showed maximum activity, while the remaining compounds have also shown moderate antibacterial activity.

QSAR Study

In order to identify substituent effect of antimicrobial activity, we established a Quantitative Structure Activity Relationship (QSAR) between in-vitro antimicrobial activity and descriptors coding for electronic properties of the molecule under consideration using linear regression analysis [17-18]. Reported antimicrobial activity data in terms of MIC values were first converted into p(MIC) values and used as a dependent variable in the QSAR study. The different molecular descriptors (independent variable) like Surface Area Approximation (SAA), Surface Area Grid (SAG), Volume (VOL), Hydration Energy (HE), Log P, Refractivity (REF), Polarizability (POL), Mass, Total Energy (TE), Electronic Energy (EE), Heat of Formation (HF), Dipole Moment (DM) and Zero Point Energy (ZPE) calculated for synthesized compounds.

The generated QSAR model was selected on the basis of various statistical parameters such as correlation coefficient which is relative measure of quality of fit, Fischer's value (F-test) which represent F-ratio between the variance of calculated and observed activity, standard error, representing absolute measure of quality of fit respectively. The best QSAR equations are given below:-

S.aureus/AM1

$$p(\text{MIC}) = -0.00013(\text{SAA}) - 0.00131(\text{VOL}) + 0.017406(\text{HE}) + 0.461515$$

$$N=12, SD = 0.071211, R = 0.79799, F = 4.675247$$




Figure 1

Table 4

pMIC (observed)	PMIC(Calculated)
-1.2	-1.23
-1.36	-1.39
-1.26	-1.27
-1.26	-1.35
-1.15	-1.27
-1.4	-1.39
-1.38	-1.35
-1.34	-1.27
-1.3	-1.24
-1.38	-1.38
-1.41	-1.32
-1.11	-1.11

Table 5

pMIC (observed)	PMIC(Calculated)
-1.2	-1.22
-1.36	-1.38
-1.26	-1.33
-1.26	-1.18
-1.15	-1.21
-1.4	-1.38
-1.38	-1.33
-1.34	-1.32
-1.3	-1.23
-1.38	-1.38
-1.41	-1.28
-1.11	-1.09










Figure 2

Table 5

Compound	Mol. formula	Mol. wt.	M. P. (°c)	Yield (%)
A-1	C ₂₀ H ₁₇ N ₇ O ₂ S	419.46	140-142	71
A-2	C ₂₁ H ₁₉ N ₇ O ₃ S	449.49	143-145	67
A-3	C ₂₀ H ₁₇ N ₇ O ₃ S	435.46	152-154	63
A-4	C ₂₂ H ₁₉ N ₇ O ₂ S	445.5	194-196	79
A-5	C ₂₁ H ₁₇ N ₆ O ₂ S	418.47	178-180	58
A-6	C ₂₂ H ₂₀ N ₆ O ₃ S	448.5	145-147	56
A-7	C ₂₁ H ₁₈ N ₆ O ₃ S	434.47	146-148	65
A-8	C ₂₃ H ₂₀ N ₆ O ₂ S	444.51	186-188	70
A-9	C ₂₁ H ₁₉ N ₅ OS	389.47	126-128	63
A-10	C ₂₂ H ₂₁ N ₅ O ₂ S	419.5	139-141	64
A-11	C ₂₁ H ₁₉ N ₅ O ₂ S	405.47	146-148	68
A-12	C ₂₃ H ₂₁ N ₅ OS	415.51	165-167	72

Table 6: Structures of the compounds under study

Code	Compound	Structure
A-1	1-(5-benzylideneamino)-3-(methylthio)-1-(pyrazine-2-carbonyl)-1H-pyrazole-4-yl)-3-methyl-1H-pyrazole-5(4H)-one	
A-2	1-(5-(4-benzylideneamino)-3-(methylthio)-1-(pyrazine-2-carbonyl)-1H-pyrazole-4-yl)-3-methyl-1H-pyrazole-5(4H)-one	

A-3	1-(5-(2-benzylideneamino)-3-(methylthio)-1-(pyrazine-2-carbonyl)-1H-pyrazole-4-yl)-3-methyl-1H-pyrazole-5(4H)-one	
A-4	3-methyl-1-(3-(methylthio)-5-(3-phenylallylideneamino)-1-(pyrazine-2-carbonyl)-1H-pyrazole-4-yl)-1H-pyrazole-5(4H)-one	
A-5	1-(5-(benzylideneamino)-1-isonicotinoyl-3-(methylthio)-1H-pyrazole-4-yl)-3-methyl-1H-pyrazole-5(4H)-one	
A-6	1-(1-isonicotinoyl-5-(4-methoxybenzylideneamino)-3-(methylthio)-1H-pyrazole-4-yl)-3-methyl-1H-pyrazole-5(4H)-one	
A-7	1-(5-(2-hydroxybenzylideneamino)-1-isonicotinoyl-3-(methylthio)-1H-pyrazole-4-yl)-3-methyl-1H-pyrazole-5(4H)-one	
A-8	1-(1-isonicotinoyl-3-(methylthio)-5-(3-phenylallylideneamino)-1H-pyrazole-4-yl)-3-methyl-1H-pyrazole-5(4H)-one	





A-9	1-(5-(benzylideneamino)-3-(methylthio)-1-phenyl-1H-pyrazole-4-yl)-1H-pyrazole-5(4H)-one	
A-10	1-(5-(4-methoxybenzylideneamino)-3-(methylthio)-1-phenyl-1H-pyrazole-4-yl)-3-methyl-1H-pyrazole-5(4H)-one	
A-11	1-(5-(2-hydroxybenzylideneamino)-3-(methylthio)-1-phenyl-1H-pyrazole-4-yl)-3-methyl-1H-pyrazole-5(4H)-one	
A-12	3-methyl-1-(3-(methylthio)-5-(3-phenylallylideneamino)-1-phenyl-1H-pyrazole-4-yl)-1H-pyrazole-5(4H)-one	

Table 7: Various computed parameters

Compd.	SAA		SAG		VOL		HE		Log P	
	AM1	PM3	AM1	PM3	AM1	PM3	AM1	PM3	AM1	PM3
A-1	542.45	544.31	666.28	672.93	1152.3	1161.82	-7.52	-7.69	-0.82	-0.82
A-2	599.46	588.85	709.75	699.53	1231.41	1211.02	-9.28	-8.83	-1.81	-1.41
A-3	526.83	530.08	671.28	673.12	1164.56	1167.43	-9.49	-9.99	-1.84	-1.84
A-4	559.03	559.2	720.27	713.79	1233.66	1228.91	-7.28	-7.47	-0.41	-0.41
A-5	552.17	552.13	677.77	680.99	1161.71	1170.39	-6.98	-7.09	-0.32	0.32
A-6	610.13	610.46	719.62	730.67	1240.43	1248.46	-8.66	-8.74	-0.68	-0.68
A-7	552.62	556	677.99	687.46	1180.24	1190.42	-10.88	-11.54	-0.71	-0.71
A-8	528.44	537.77	683.14	696.56	1202.3	1219.59	-5.43	-5.5	-0.66	-0.66
A-9	564.69	567.71	674.02	684.52	1172.57	1181.96	-5.31	-5.43	1.79	1.79
A-10	622.33	624.01	723.77	735.72	1251.39	1258.79	-7.02	-7.09	0.8	0.8
A-11	552.64	554.53	680.73	695.22	1182.47	1194.59	-9.33	-9.31	0.76	0.76
A-12	374.81	542.62	657.73	717.88	1139.15	1243.63	-2.1	-4.83	2.2	2.2

Table 8: Various computed parameters (cont'd)

Compd.	RF		POL		Mass		TE		EE	
	AM1	PM3	AM1	PM3	AM1	PM3	AM1	PM3	AM1	PM3
A-1	120.24	120.24	43.96	43.96	419.46	419.46	-116722	-104613	-933098	-891930
A-2	126.62	126.52	46.43	46.43	449.43	449.49	-127947	-114833	-1E+06	-1E+06
A-3	121.85	121.85	44.6	44.6	435.46	435.46	-124364	-111391	-986335	-960788
A-4	130.49	130.49	47.44	47.44	445.5	445.5	-123459	-110732	-999302	-981066
A-5	122.39	122.39	44.67	44.67	418.47	418.47	-115478	-103965	-912033	-892043
A-6	128.76	128.76	47.14	47.14	448.5	448.5	-126452	-114179	-1E+06	-994683
A-7	123.99	123.99	45.31	45.31	434.47	434.47	-122870	-110741	-976927	-955952
A-8	132.65	132.65	47.85	47.85	447.51	447.51	-124879	-113471	-1E+06	-1E+06
A-9	125.96	125.96	45.38	45.38	417.49	417.49	-113980	-103314	-909874	-888750
A-10	132.33	132.33	47.85	47.85	447.51	447.51	-124955	-113528	-1E+06	-992207
A-11	127.56	127.56	46.02	46.02	433.48	433.48	-121106	-110093	-1E+06	-960111
A-12	136.2	136.2	48.86	48.86	443.52	443.52	-132217	-109384	-1E+06	-962773

Table 9: Various computed parameters (cont'd)

Compd.	HF		DM		ZPE	
	AM1	PM3	AM1	PM3	AM1	PM3
A-1	423.45	112.18	1.77	0.615	226.12	216.56
A-2	134	68.03	1.57	2.07	246.98	237.55
A-3	130.81	67.02	1.82	2.18	228.83	219.92
A-4	229.87	172.52	3.05	3.58	246.72	238.14
A-5	158.6	102.15	2.25	2.2	233.82	224.29
A-6	120.15	63.13	3.35	3.17	254.72	244.37
A-7	115.61	58.59	3.47	3.59	236.67	227.72
A-8	184.81	113.49	4.62	4.5	256.33	247.46
A-9	147.45	95.18	0.52	1.18	241.45	232.01
A-10	108.59	68.28	1.07	0.51	262.36	252.09
A-11	370.42	47.7	2.07	2.1	284.24	236.1
A-12	474.93	203.85	2.45	8.76	330.07	251.48

4. Conclusion

The present study involves some pyrazoles compounds (A-1 to A-12) to check their QSAR studies pertaining to their antibacterial activity against *S.aureus*. This may be concluded from the studies that the compounds A-2, A-3, A-6, A-10 and A-11 showed significant antibacterial activity against *S.aureus*. The present paper also discusses Quantitative Structure Activity Relationship equation. In conclusion this may be concluded that following parameters shows positive contributions towards p(MIC).

AM1/ *S.aureus*:- SAA, VOL, HE

PM3/ *S.aureus*:-HE, TE, HF

5. References

1. Udupi R.H., Narayan Rao S., Bhat A.R., "Synthesis of some new pyrazolone derivatives as anti-microbial, anti-inflammatory & analgesic agents", Ind. J. Het. Chem., **1998**, *7*, 217.
2. Halnor V.B., Joshi N.S., Karale B.K., Gill C.H., "Synthesis & biological activities of some pyrazolones", Ind. J. Het. Chem., 2005, *14*, 371.
3. Bondock S., Fadaly W., Metwally M.A., "Synthesis and antimicrobial activity of some new thiazole, thiophene and pyrazole derivatives containing benzothiazole moiety", Eur. J. Med. Chem., **2010**, *45*, 3692.
4. Radi S., Salhi S., Radi A., "Synthesis and preliminary biological activity of some new pyrazole derivatives as acyclonucleoside analogues", Letters in Drug Design & Discovery, **2010**, *7*, 27.
5. Sahu S.K., Banerjee M., Samantra A., Behera C., Azam M.A., "Synthesis, Analgesic, anti-inflammatory and antimicrobial activities of some novel pyrazolone derivatives", Tropical J. Pharm. Research, **2008**, *7*(2), 961.
6. Sridhar S., Perumal P.J., Etti S., Shanmugam G., Ponnuswamy M.N., Prabavathy V.R., Mathivanan N., "Design, synthesis and antimicrobial activity of 1H-pyrazole carboxylate", Bioorg. Med. Chem. Lett., **2004**, *14*, 6035.
7. Rao Jyothi N., Sujith K.V., Kalluraya B., "An efficient microwave assisted synthesis of some novel pyrazolone and their biological activity", J. Saudi Chem. Soc., 2008, *12*(3): 347.
8. Satheesha Rai N., Balakrishna Kalluraya, "A novel synthesis of nitro furan containing 1, 3, 4, 5-tetra substituted pyrazoles via 1, 3-dipolar addition reaction", Indian J. Chem., **2007**, *46B*, 375.
9. Argade N.D., Kalrale B.K., Gill C.H., "Microwave assisted improved method for the synthesis of

- pyrazole containing 2, 4-disubstituted oxazole-5-one and their antimicrobial activity”, *E-Journal Chemistry*, **2007**, 5(1), 120.
10. Chovatia P.T., Akabari J.D., Kachhadia P.K., Zalavadia P.D., Joshi H.S., “Synthesis and selective antitubercular and antimicrobial inhibitory activity of 1-acetyl-3, 5-diphenyl-4, 5-dihydro-(1H)-pyrazole derivatives”, *J. Serb. Chem. Soc.*, **2007**, 71(7): 713.
 11. Kuntal M., Yadvendra K., “Microwave assisted synthesis of new indophenazine 1, 3, 5-trisubstituted pyrazolone derivatives of benzofuran and their antimicrobial activity”, *Bioorganic and Medicinal Chemistry Letters*, **2009**, 19: 2688.
 12. Karthikeyan M.S., Holla B.S., Kumari N.S., “Synthesis and antimicrobial studies on novel chloro fluorine containing hydroxyl pyrazolones”, *Eur. J. Med. Chem.*, **2007**, 42, 30.
 13. Venkat Raghvan R., Vijayakumar V., Suchethakumari N., “Synthesis and antimicrobial activities of novel 1, 5-diaryl pyrazole”, *Eur. J. Med. Chem.*, **2010**, 45: 1173.
 14. Menozzi G., Merello L., Fossa P., Schenone S., Ranise A., Mosti et al, “Synthesis, antimicrobial activity and molecular modeling studies of halogenated 4-[1H-imidazole-1-yl(phenyl) methyl]-1, 5-diphenyl-1H-pyrazole”, *Bioorganic and Medicinal Chemistry*, **2004**, 12: 5465.
 15. Pattan R., Rabara P.A., Pattan J.S., Bukitagar A.A., Wakale V.S., Musmade D.S., “Synthesis and evaluation of some new substituted phenyl thiazole derivatives and their anti-tubercular activity”, *Indian Journal of Chemistry*, **2009**, 48B: 1453.
 16. Arora K., Nathani V., “Antimicrobial and QSAR studies of some pyrazolones compounds”, *Research Journal of Pharmaceutical Biological and Chemical Sciences*, **2012**, 3(4): 1423.
 17. Arora K., “3D QSAR studies for some pharmacological important compounds”, *International Journal of Pharma and Biosciences*, **2014**, 5(1): 571-579.