



Research Article
**International Journal of Pharmacy and
Natural Medicines**

www.pharmaresearchlibrary.com/ijpnm



**Preliminary Phytochemical Investigation, Antiproliferative Activity and
Antioxidant Activity of extracts of *Fritillaria roylei***

Tufael Amin Mir and Wajaht A. Shah

Department of Chemistry, University of Kashmir, Hazratbal Srinagar, India-190006
Received: 17 March 2014, Accepted: 24 May 2014, Published Online: 15 June 2014

Abstract

Fritillaria roylei is one of the medicinal plants used in traditional medical system for ailment of various diseases. But scientific study of the plant has been less carried out for which the study was conducted for validation of prevailing medical practice. The Soxhlet extraction of the plant were conducted in various solvents (hexane, chloroform, ethyl acetate, methanol and water) differing in polarity. The phytochemical screening of extracts revealed the presence of alkaloids, saponins and tanins. In vitro Cytotoxicity evaluation against five human cancer cell lines A549 (Lung), C6 (Glioma), T47D (Breast), MCF (Breast), and TH-1 (Colon) showed that the extracts of *Fritillaria roylei* exhibited potent cytotoxic activity. The extracts of *Fritillaria roylei* also showed antioxidant activity.

Keywords: *Fritillaria roylei*, cytotoxic, antioxidant

Contents

| | |
|------------------------------------|----|
| 1. Introduction | 94 |
| 2. Materials and Method. | 95 |
| 3. Results and Discussion. | 96 |
| 4. References | 97 |

***Corresponding author**

Wajaht A. Shah

University of Kashmir,
Hazratbal Srinagar, India -190006
E-mail: doctorwajaht@gmail.com
Manuscript ID: IJPNM2038



PAPER-QR CODE

Copyright © 2013, IJPNM All Rights Reserved

1. Introduction

Nature stands a golden mark and provides the storehouse of remedies to cure all ailments of mankind. Herbs have always been the principal form of medicine throughout the world as people strive to stay healthy in the face of chronic stress and pollution and to treat illness with medicines that work in count with the body's own defense. There is a widespread belief that green medicines are healthier and safer than synthetic ones (Sindhia & Bairwa 2010). In traditional medicine, there are many natural crude drugs that have the potential to treat many diseases and disorders and one of them is *Fritillaria roylei*. *Fritillaria* is a genus of about 1000 species of bulbous plants belonging to the family Liliaceae. It is a perennial temperate herb that grows in mountain slopes and sub-alpine meadows, usually on open, stony and moist hill sides. *F. roylei* is a rare species found at an altitude around 3800- 4700 m. In the West, *Fritillaria* is most commonly regarded as an ornamental garden plant. It is traditionally valued as an herbal remedy

in Nepal and China. *Fritillaria* affects the heart and lung meridians orenergy pathways in the body and is used primarily to treat various lung diseases, including asthma, bronchitis, tuberculosis and coughs of any type. It is also used as a lymphatic decongestant to reduce swellings, nodules, fibrocystic breasts, goiter and swollen lymph glands. Previous research reveals that it is used for treatment of nervous system, prolonged decrease in blood pressure, stimulation of the heart muscle and dysfunction of breathing (Erika & Rebecca 2005). Plant contains phytochemicals to protect itself. But recent researches demonstrate that many phytochemicals can protect human and plants diseases (Lata & Dubey 2010). Compounds theorized to be responsible for activity includes several bioactive isosteroidal alkaloids (verticine, verticinone, isovericine, imperialine, hupe henine, ebeiedine, ebeienine and ebeiedinone) and 2 nucleosides (thymidine and adenosine) (Erika & Rebecca 2005). *Fritillaria roylei* possess peimine, peiminine, peimisine, peimiphine, peimidine, peimitidine, propeimin, sterol, etc and has been reported for antiasthmatic, antirheumatic, febrifuge, galactagogue, haemostatic, ophthalmic and oxytocic properties, etc (Chi, 1936; Wu, 1944; Jiang, 2001). Very less pharmacological and anticancer study has been reported on *F. roylei* for which the study was designed to establish scientific basis for some of its therapeutic properties in traditional use. This investigation encourages for further findings of medicinal plants. Hence, the study deals with the phytochemical screening, cytotoxic assay and antioxidant activity that helps in scientific evaluation of bioactive compounds and may prove beneficial for mankind.

2. Materials and Method

Methodology

Collection of samples

The plant material were collected from Gulmarg area in the month of May- June 2013. They were shade dried and pulverized in powder form. Voucher specimen of *F. roylei* bearing specimen no.27913, was deposited at KASH herbarium in centre of plant taxonomy, Kashmir University, Srinagar

Preparation of extract

The powdered sample were extracted by soxhlet extraction method using hexane, chloroform, ethyl acetate, methanol and water as solvent in the order of increasing polarity. Then, solvents from extracts were removed under reduced pressure with the aid of rotary vacuum evaporator.

Experimental

Phytochemical Screening of Crude Extracts.

The phytochemical screening of the plant extracts was carried out according to standard procedure (Farnsworth et al, 1966)

Alkaloids:

The methanolic extract (30ml) was evaporated to dryness in an evaporating dish on water bath. Five ml of 2 N HCl were added and stirred while heating on the water bath for 10min., cooled, filtered, and the filtrate was treated with a few drops of Mayer reagent. The samples were then observed for the presence of turbidity or precipitation.

Flavonoids:

The alcoholic extract (75 ml) of plant sample were evaporated to dryness on a water bath, cooled and the residue was defatted by washing several times with petroleum ether. The defatted residue was dissolved in 30ml 80% ethanol and filtered. The filtrate was treated with a few drops of concentrated HCl and magnesium turnings (0.5g). The presence of flavonoids was indicative if a pink or magenta red color developed within 3 min.

Tannins:

The alcoholic extract (25ml) was evaporated to dryness on a water bath. The residue was extracted several times with n-hexane & filtered, the insoluble residue was stirred with 10ml of hot saline solution, the mixture was cooled, filtered and the volume of filtrate was adjusted to 10ml with more saline solution. To 5 ml of this solution, few drops of ferric chloride test reagent were added. An intense green blue or black colour was taken as an evidence for the presence of tannins.

Saponins: 1 gm of ethanol extract was dissolved in 10ml of distilled water in a test tube and shaken vigorously for 1-2 min. The presence of saponins was indicated by characteristic honeycomb froth at least 1cm in height, which persisted for 30 min

Anticancer activity

Human cell lines and culture

Cytotoxic assay was carried out by MTT (3-(4,5-Dimethylthiazol-2-yl)- 2,5 diphenyltetrazolium bromide, a tetrazole) protocol in order to evaluate the anti proliferative effect of extracts of *F. roylei*. For this purpose, a sufficient number of exponentially growing cells were used to avoid confluence of the culture during the treatment. The cell lines A549, C6, T47D, MCF, and TH-1 were seeded at 10^4 cells/well and allowed to adhere for 12 h.

Cytotoxicity assay

In order to evaluate the optimum concentration at which the extracts inhibited the cell proliferation in all the five cell lines, cells were treated with the extracts at a concentration of 100 µg/ml. DMSO was used as a solvent for the

dilution of extract, which was also used as an experimental control. Mitomycin-C was used as positive controls at a concentration of 1×10^{-5} $\mu\text{g/ml}$. After 48 h treatment, cell growth was evaluated by MTT assay (Alley et al. 1986, 1988). MTT solution of 50 μl (5 mg/ml of PBS) was added to each well and the plates were incubated for 3 h at 37°C in dark. The media was aspirated and 150 μl of MTT solvent (4 mM HCl, 0.1% Nondet P-40, all in isopropanol) was added to each well to solubilize the formazan crystals. The absorbances of plates were measured on ELISA reader (Benchmark, BioRad) at a wavelength of 570 nm. Each sample was performed in triplicate, and the entire experiment was repeated thrice

Antioxidant Activity:

The free radical scavenging activity of the extract was measured in terms of hydrogen donating or radical-scavenging ability using the stable radical DPPH. 0.1 mM solution of DPPH in methanol was prepared and 1.0 ml of this solution was added to 1.0 ml of the test solution in methanol at concentrations of 100 $\mu\text{g/ml}$. Thirty minutes later, the absorbance was measured at 517 nm. Ascorbic acid was used as the reference compound. Lower absorbance of activity was expressed as the inhibition percentage of free radical by the sample and was calculated using the following formula:

$$\% \text{ scavenging activity} = (A_0 - A) / A_0 \times 100.$$

Where A_0 was the absorbance of the control (blank, without compound) and A was the absorbance of the the reaction mixture. All the tests were performed in triplicate and the graph was plotted with the mean values.

3. Results and Discussion

The use of herbs for medicinal purpose is a universal phenomenon. Every culture and civilization on earth, through written or oral tradition, has relied on vast variety of natural chemistries found in plants for their therapeutic properties. All drugs from the plant are substances with a particular therapeutic action extracted from plants. Plant substances continue to serve as viable source of drugs for the world population and several plant-based drugs are in extensive clinical use (Rosa & Rao 2000). The curative properties of plants are perhaps due to the presence of various secondary metabolites which are non nutritive plant compounds. These classes (such as alkaloids, tannins, anthraquinones and flavonoids) of compounds are known to have curative activity against several pathogens and therefore could suggest the use of it traditionally for treatment of various illnesses (Usman & Osuji 2007). The extracts of *Fritillaria roylei* were phytochemically screened for the presence of Alkaloids, Flavonoids, Tannins and Saponins. However the results showed that the *Fritillaria roylei* contains Alkaloids, Tannins and Saponins in good amount. Since these alkaloids and tannins are responsible for broad pharmacological profile, including antiinflammation, antioxidant, antibacterial, anticancer, and anti-histamine activity.

Cytotoxic activity:

Therefore the methanolic and ethylacetate extracts of the *Fritillaria roylei* were screened for their possible anticancer activity. In order to understand the effects of various extracts of *Fritillaria roylei* on human cancer cell lines, experiments were carried using cultured A549 (lung), C6 (Glioma), T47D (Breast), MCF (Breast) and TH-1 (Colon) cell lines by MTT assay. All cell lines were submitted to 100 μg concentration of extracts, which reduced the viability of these cell lines. As shown in Table 1 the extract was active against all the five cancer cell lines tested. The concentration of Mitomycin C was taken as 1×10^{-6} $\mu\text{g/ml}$.

Table 1. In-vitro cytotoxicity of the extracts of *Fritillaria roylei*

| Cell-type | A549 | C6 | T47D | MCF | TH-1 |
|----------------------|------|----|------|-----|------|
| DMSO | 6 | 6 | 4.5 | 6.5 | 7 |
| Mitomycin C | 92 | 89 | 91 | 78 | 83 |
| Methanolic extract | 65 | 71 | 53 | 62 | 76 |
| EthylAcetate extract | 55 | 43 | 77 | 78 | 57 |

Antioxidant Activity:

DPPH free radical scavenging capacity of methanolic extract of *Fritillaria roylei* was measured by DPPH assay under in vitro conditions. The ability of the examined extract to act as donor for hydrogen atoms in the transformation of DPPH radical into its reduced form DPPH₂ was investigated. The examined samples were able to reduce the stable purple colored DPPH radical into yellow colored DPPH₂. The more rapidly the absorbance decreases, the more potent the antioxidant activity of the sample in terms of its hydrogen atom-donating capacity (Shahidi, Alasalvar, & Liyana-Pathirana, 2007; Alasalvar et al., 2009). The aforementioned extract showed most promising radical scavenging activity at concentration of 150 $\mu\text{g/ml}$. The results are plotted in the form of graph (fig.1).




Figure 1. Antioxidant activity of methanolic extract of *Fritillaria roylei*

4. References

1. Alley, M.C., Scudiere, D.A., Monks, A., Hursey, M.L., Czerwinski, M.J., Fine, D.L., Abbott, B.J., Mayo, J.G., Shoemaker, R.H., Boyd, M.R., Feasibility of drug screening with panels of human tumor cell lines using a micro culture tetrazolium assay. *Cancer Res.*, **1988**, 48: 589–601.
2. Alasalvar, C., Magdalena, K., Agnieszka, K., Rybarczyk, A., Shahidi, F., & Amarowicz, R., Antioxidant activity of hazelnut skin phenolics. *Journal of Agricultural and Food Chemistry*, **2009**, 57: 4645–4650
3. Chi Yuoh-Fong, Kao Y S. and Chang K.J., The Alkaloids of *Fritillaria Roylei*. Isolation of Peimine1, *J. Am. Chem. Soc.*, **1936**, **58**: 1306–1307.
4. Erika, L and J. F. Rebecca. 2005. *Gale encyclopedia of publichealth*. Gale Cengage Learning. Gonzalez, A., G. Maujir., I. Bazzocchi., I. L. Correa and M.D. Gupta. Screening of antimicrobial and cytotoxicactivities of Panamanian plants. *Phytomedicine* 1, **1994**: 149-153.
5. Farnsworth NR Biological and phytochemical screening of plants. *J. Pharm. Sci.*, **1966**, 55: 225-276.
6. Lata, N. and V. Dubey., Preliminary phytochemical screening of *Eichhornia crassipes*: the world's worst aquatic weed. *J. of Pharmacy Research*, **2010**, 3(6): 1240-1242
7. Rosa, G. and P. S. Rao., Anticancer compounds from tissue cultures of medicinal plant. *J. of Herbs and Spices Med. Plants*, **2000**, **7**: 71-102.
8. Shahidi, F., Alasalvar, C., & Liyana-Pathirana, C. M. **2007**., Antioxidant phytochemicals in Hazelnut
9. kernel (*Corylus avellana* L.) and Hazelnut Byproducts. *Journal of Agricultural and Food Chemistry*, 55, 1212–1220.
10. Sindhia, V. R. and R. Bairwa, Plant review: *Buteamonosperma*. *Int. J. of Pharmaceutical and Clinical Research*, **2010**, **2**: 90-94.
11. Usman, H. and J. C. Osuji., Phytochemical and invitro antimicrobial assay of the leaf extract of *Newbouldia leavis*. *Afr. J. Trad. CAM*, **2007**, 4(4): 476-480.