



International Journal of Chemistry and Pharmaceutical Sciences

Journal Home Page: www.pharmaresearchlibrary.com/ijcps



Research Article

Open Access

Exploiting the Antiparasitic Activity of Naphthalimides Derivatives

Gabriela Cristo^{1,2}, Luís Gaspar^{1,2}, Jennifer Noro³, Catarina Baptista^{1,2}, Paul Kong Thoo-Lin⁴, Maria José Alves³ and Anabela Cordeiro-da-Silva^{1,2,5,*}

¹Instituto de Investigação e Inovação em Saúde, Universidade do Porto, Porto, Portugal

²IBMC-Institute for Molecular and Cell Biology, Parasite Disease Group, Porto, Portugal

³Department of Chemistry, Campus de Gualtar, University of Minho, Braga, Portugal;

⁴School of Pharmacy and Life Sciences, Robert Gordon University, Riverside East, Garthdee Road, Aberdeen AB101GJ, Scotland

⁵Department of Biological Sciences, Faculty of Pharmacy, University of Porto, Porto, Portugal

ABSTRACT

A set of 1,8-naphthalimides derivatives were synthesized and tested against three protozoans that cause important human diseases: *Leishmania infantum*, *Trypanosomabrucei* and *Trypanosoma cruzi*. Additionally, toxicity was determined by growth inhibition of THP-1 derived macrophages. The results suggest that chemical modifications in the carbon chain linking the naphthalimide and the substituting groups have different effects in the parasites. This work should provide new insights for the design and optimization of more potent and directed naphthalimide derivatives against these organisms.

Keywords: *Leishmania infantum*, *Trypanosoma brucei*, *Trypanosoma cruzi*

ARTICLE INFO

CONTENTS

1. Introduction	19
2. Experimental	20
3. Results and Discussion	20
4. Acknowledgement	22
5. References	22

Article History: Received 18 November 2015, Accepted 26 December 2015, Available Online 27 January 2016

*Corresponding Author

Anabela Cordeiro-da-Silva
Instituto de Investigação e Inovação em Saúde,
Universidade do Porto, Porto, Portugal,
Manuscript ID: IJCPS2793



PAPER-QR CODE

Citation: Anabela Cordeiro-da-Silva, et al. Exploiting the Antiparasitic Activity of Naphthalimides Derivatives. *Int. J. Chem, Pharm, Sci.*, 2016, 4(1): 19-23.

Copyright© 2016 Anabela Cordeiro-da-Silva, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original work is properly cited.

1. Introduction

Parasitic diseases caused by trypanosomatids are still an important health problem, mainly in tropical and subtropical areas. In addition, temperate regions of our globe, including North America and the Asia-Pacific region

International Journal of Chemistry and Pharmaceutical Sciences

are also affected by disease-causing protozoans like *Leishmania spp.*, *Trypanosoma cruzi* and *Trypanosoma brucei*. These are the species of trypanosomatids most associated with human health, being responsible for high mortality and morbidity[1]. There are still no vaccines and the actual chemotherapies are far from satisfactory, owing to the emergence of resistances, serious side effects, and its limited efficacy. Therefore, it is imperative to continue the discovery of new drugs to treat these diseases[2].

2. Experimental

Naphthalimides and bis-naphthalimides are classes of compounds bearing aromatic groups that have generated intense interest by scientists around the world because of their many reported biological activities. Special attention has been devoted to the high anticancer activity of naphthalimides, which is due to their interactions with

DNA by a mechanism of intercalation[3–5]. In addition, recent studies showed that 1,8-naphthalimide derivatives also demonstrated other biological activities, such as antitrypanosomal [6]. In this context, this work proposes to study the relationship between the structure and the activity of a set of 1,8-naphthalimide derivatives with 2, 3 or 4 carbons linking the naphthalimide moiety to different functional groups (amine, imine, guanidine, urea, amide, and 1,2,3-triazole) against three trypanosomatids: *Leishmania infantum*, *Trypanosoma brucei* and *Trypanosoma cruzi*. These 1,8-naphthalimide derivatives compounds had been recently synthesized and include naphthalimidoalkyl amines 1a, b and its heterocyclic imine derivatives 2a-l, heterocyclic amine derivatives 3f,g, guanidine derivatives 4a-c, urea derivatives 5a-l, amide derivatives 6a-l and triazol derivatives 7d-f[7,8].




Figure 1: Naphthalimides compounds 1a,b, 2a-l, 3f,g, 4a-c, 5a-l, 6a-l, 7d-f used in trypanosomatids tests: *Leishmania infantum*, *Trypanosoma brucei* and *Trypanosoma cruzi*

3. Results and Discussion

All the naphthalimides compounds stock solutions were prepared in DMSO in a concentration of 10 mm and stored at -20 °C. Anti *T. cruzi* (Y strain) activity was performed by high content screening (adapted from [9]); for *T. brucei* (L427 Wild Type) blood stream forms and axenic amastigote of *L. infantum* (clone MHOM/MA671TMA-P263) activity was determined using the resazurin-based assay[10,11]; intracellular amastigote activity of *L. infantum* was performed by luciferase assay (adapted from[12]). The cellular toxicity of the compounds was evaluated on THP-1 differentiated macrophages using the MTT assay[13]. The table summarizes the results obtained for the compounds with relevant anti-parasitic activity at 10 μM concentration. Overall, our results show that the best activity was against

T. brucei. The group of naphthalimidoalkyl amines represented by compound 1a and 1b show strong inhibition at concentration 10 μM (101±1% inhibition); however compound 1a presents higher toxicity with NOAEL (no observed adverse effect level) > 25 μM in comparison with 1b NOAEL > 100 μM, suggesting that shorter a carbon chain (n =2) increases the cytotoxicity. The group of heterocyclic imines compounds 2a-l generally reveals good activity against *T. brucei*. Within the imino furan subgroup, compounds 2b, 2f and 2j, only 2f with 3 carbon atoms chain shows to be active (101 ± 1% inhibition), and in the subgroup incorporating imidazole (2d, 2h and 2l) compound 2l, with 4 carbon atoms chain, is the least active. The toxicity for heterocyclic imines is lower with 3 and 4 carbon chains. The compound with the best balance activity-toxicity belongs to

compound 2e, with a three-carbon atom chain and a pyrrole unit. Heterocyclic amines compounds are also active against *T. brucei*, our results showing almost complete inhibition with compound 3f (99 ± 1% inhibition) and 3g (96 ± 3% inhibition). However, compound 3g shows lower cytotoxicity than 3f, with NOAEL > 50µM and > 25µM, respectively. Guanidine substituted compounds 4a-also reveals some anti-*T. brucei* activity, with a clear tendency for toxicity to decrease with the increase of the carbon chain length. Within the urea-substituted compounds 5a-l the most active compounds against *T. brucei*, at the concentrations tested, is 5f and 5l both with 4 carbon atom chains. Within this group, when the carbon chain increases up to four carbon atoms, so does the activity increase. Also the nature of the terminal groups in compounds 5a-lis also found to be important for activity. Comparing the cytotoxicity of the four carbon chain members in this group (5c, 5f, 5i and 5l), only compound 5c was found non toxic, suggesting that un-substituted ureas are less toxic than substituted ureas against THP-1 cell line. In relation to the triazole group, only compound 7f shows activity against *T. brucei* (90 ± 3% inhibition), with toxicity increasing when the triazole group is linked by a 3 carbon atoms chain. Regarding *T. cruzi*, amine 1a and guanidine 4 shows modest anti-parasitic activity (34 ± 9% inhibition and 45 ± 9% inhibition, respectively). Urea 5l bearing a 4

carbon atoms chain and a *n*-butyl group as substituent shows to be the most active compound of all tested against *T. cruzi*, (81 ± 8% inhibition). However, all compounds that show some activity against *T. cruzi* are accompanied by an increase in cytotoxicity, suggesting a nonspecific mechanism. Previous studies conducted in our laboratory have shown that bis-naphthalimidopropyl (BNIP) derivatives compounds exert significant effects against *L. infantum* [14]. Surprisingly, no significant anti-leishmanial activity was found for all the groups studied; either against *L. infantum* intracellular and axenic amastigotes. A hypothesis is that the activity is dependent on a second naphthalimide group in the molecule. Overall, the toxicity of the synthesized compounds with a 2 carbon atoms linker presents the higher toxicity. An exception was observed with the substituted urea-derived group, where the 4 carbon atoms chain compounds are the most toxic, with the exception urea 5c. The group of amides 6a-l compounds does not present relevant anti-parasitic activity (data not shown). In conclusion, naphthalimides bearing primary alkylamines and un-substituted ureas are the hits for further research in *Trypanosoma brucei*, in combination with longer linkers, and *O*-, *N*-functionalized linkers. No interesting results have been found to the other two trypanosomatids studied.

Table 1: Anti-parasitic activity against *Leishmania infantum* intracellular amastigotes and axenic amastigotes, *Trypanosoma brucei*, *Trypanosoma cruzi* and cytotoxicity in THP1-derived macrophages for the compounds tested.

Functional groups: naphthalimidoalkylamines		10µM single dose testing (%activity ± SD)				Toxicity NOAEL (µM)
Compound	Radical / Number of carbons	<i>Leishmania infantum</i> intracellular amastigotes	<i>Leishmania infantum</i> axenic amastigotes	<i>Trypanosoma brucei</i>	<i>Trypanosoma cruzi</i>	
1a	R=NH ₂ / n=2	23 ± 17	5 ± 5	101 ± 1	34 ± 9	> 25
1b	R=NH ₂ / n=3	N. A.	N. A.	101 ± 1	N. A.	> 100
Functional group: Heterocyclic imine						
2a	X=NH/Y=C/ n=2	N. A.	9 ± 7	101 ± 2	22 ± 13	> 25
2b	X=O/ Y=C/ n=2	N. A.	5 ± 0	17 ± 12	N. A.	> 25
2c	X=S/Y=C/ n=2	N. A.	1 ± 3	8 ± 5	N. A.	> 25
2d	X=NH/Y=N/ n=2	N. A.	2 ± 1	101 ± 1	25 ± 4	> 25
2e	X=NH/Y=C/ n=3	N. A.	N. A.	101 ± 1	N. A.	> 100
2f	X=O/Y=C/ n=3	N. A.	20 ± 21	101 ± 1	N. A.	> 50
2g	X=S/Y=C/ n=3	N. A.	N. A.	101 ± 1	N. A.	> 50
2h	X=NH/Y=N/n=3	6 ± 13	23 ± 16	99 ± 2	N. A.	> 50
2i	X=NH/Y=C/ n=4	4 ± 12	1 ± 12	53 ± 5	N. A.	> 50
2j	X=O/Y=C/ n=4	N. A.	19 ± 4	26 ± 10	N. A.	> 50
2k	X=S/Y=C/ n=4	N. A.	27 ± 3	94 ± 2	N. A.	> 50
2l	X=NH/Y=N/ n=4	N. A.	24 ± 12	24 ± 3	N. A.	> 100
Functional group: Heterocyclic amine						
3f	X=O/Y=C/N=3	N. A.	13 ± 1	99 ± 1	17 ± 15	> 25
3g	X=S/ Y=C/N=3	N. A.	9 ± 4	96 ± 3	25 ± 10	> 50
Functional group: Guanidine						
4a	R=H/ n=2	N. A.	4 ± 10	100 ± 1	45 ± 9	> 25
4b	R=H/ n=3	N. A.	22 ± 8	101 ± 1	N. A.	> 50
4c	R=H/ n=4	N. A.	10 ± 2	N. A.	N. A.	> 100

Functional group: Ureas						
5a	R=H/ n=2	N. A.	14 ± 1 5	N. A.	N. A.	> 25
5b	R=H/ n=3	N. A.	27 ± 1 2	15 ± 16	3 ± 8	> 100
5c	R=H/ n=4	N. A.	16 ± 5	75 ± 12	N. A.	> 100
5d	R=4-NO ₂ C ₆ H ₄ / n=2	N. A.	11 ± 17	37 ± 8	N. A.	> 25
5e	R=4-NO ₂ C ₆ H ₄ / n=3	N. A.	N. A.	38 ± 6	26 ± 1	> 25
5f	R=4-NO ₂ C ₆ H ₄ / n=4	N. A.	10 ± 10	94 ± 4	N. A.	> 10
5g	R=CH ₂ Ph/ n=2	N. A.	30 ± 6	5 ± 4	N. A.	> 25
5h	R=CH ₂ Ph/ n=3	N. A.	16 ± 0	26 ± 6	21 ± 17	> 10
5i	R=CH ₂ Ph/ n=4	N. A.	12 ± 12	74 ± 2	5 ± 8	> 10
5j	R=(CH ₂) ₃ CH ₃ / n=2	N. A.	14 ± 1	18 ± 10	N. A.	> 50
5k	R=(CH ₂) ₃ CH ₃ / n=3	N. A.	17 ± 14	18 ± 3	20 ± 12	> 25
5l	R=(CH ₂) ₃ CH ₃ / n=4	N. A.	12 ± 12	96 ± 2	81 ± 8	> 10
Functional group: 1,2,3-Triazole						
7e	n=2	6 ± 16	13 ± 2	12 ± 4	6 ± 12	> 100
7d	n=3	N. A.	11 ± 4	16 ± 6	N. A.	> 50
7f	n=4	17 ± 2	N. A.	90 ± 3	22 ± 5	> 100

Results show means activities of at least three independent assays.

NOAEL= No-Observed Adverse Effect Level (MTT assay in PMA-differentiated THP-1 cells);

N. A. = no activity.

4. Acknowledgements

The research leading to these results has received funding from the European Community's Seventh Framework Programme under grant agreements No.602773 (Project KINDRED). L.G. was supported by the Fundação para a Ciência e Tecnologia through grant SFRH/BD/ 81604/2011. The research leading to these results has received funding from the European Community's Seventh Framework Programme under grant agreements No.602773 (Project KINDRED). Thanks are due also to the NMR Portuguese network (PTNMR, Bruker Avance III 400-Univ. Minho), and FCT and FEDER for financial support to CQU/UM.

5. References

- [1] R. Lozano, M. Naghavi, K. Foreman, S. Lim, K. Shibuya, V. Aboyans, et al. Global and regional mortality from 235 causes of death for 20 age groups in 1990 and 2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet.*, **2012**, 380, 9859: 2095–128.
- [2] L. Monzote, A. Siddiq. Drug development to protozoan diseases. *Open Med. Chem. J.*, **2011**, 5, 1–3.
- [3] M. Brana, A. Ramos. Naphthalimides as Anticancer Agents: Synthesis and Biological Activity. *Curr. Med. Chem. Agents.*, **2001**, 1: 237–255.
- [4] Z. Li, Q. Yang, X. Qian. Novel heterocyclic family of phenyl naphthothiazole carboxamides derived from naphthalimides: synthesis, antitumor evaluation, and DNA photocleavage. *Bioorg. Med. Chem.*, **2005**, 13: 3149–55.
- [5] R. Martínez, L. Chacón-García. The search of DNA-intercalators as antitumoral drugs: what it worked and what did not work. *Curr. Med. Chem.*, **2005**, 12: 127–51.
- [6] M. Muth, V. Hoerr, M. Glaser, A. Ponte-Sucré, H. Moll, A. Stich, U. Holzgrabe. Antitrypanosomal activity of quaternary naphthalimide derivatives. *Bioorg. Med. Chem. Lett.*, **2007**, 17: 1590–3.
- [7] J. Noro, J. Maciel, D. Duarte, A.C.D. Olival, C. Baptista, A.C.D. Silva. Evaluation of New Naphthalimides as Potential Anticancer Agents against Breast Cancer MCF-7, Pancreatic Cancer BxPC-3 and Colon Cancer HCT- 15 Cell Lines. *Organic Chem. Curr. Res.*, **2015**, 4, 2, 4: 144.
- [8] Compound **2h**: white solid (84.7 %); m.p.: 212–213 °C; ¹H(400 MHz, DMSO) 1.97-2.04 (m, 2H), 3.65 (t, *J*= 6.8 Hz, 2H), 4.11 (t, *J*= 7.6 Hz, 2H), 7.03 (s, 1H), 7.14 (s, 1H), 7.83 (t, *J*= 7.6 Hz, 2H), 8.19 (s, 1H), 8.41 (d, *J*= 8.4 Hz, 2H), 8.45 (d, *J*= 7.2 Hz, 2H), 12.4 (br s, 1H) ppm. Compound **2j**: brown solid (45.4 %); m.p.: 110-111 °C; ¹H(400 MHz, CDCl₃) 1.82-1.87 (m, 4H), 3.68 (t, *J*= 6.8 Hz, 2H), 4.25 (t, *J*= 7.2 Hz, 2H), 6.47 (dd, *J*= 3.2, 1.6 Hz, 1H), 6.79 (br s, 1H), 7.51 (s, 1H), 7.76 (dd, *J*= 8.4, 7.2 Hz, 2H), 8.12 (s, 1H), 8.21 (dd, *J*= 8.4, 1.2 Hz, 1H), 8.60 (dd, *J*= 7.2, 1.2 Hz, 2H) ppm. Compound **2l**: white solid (92.3 %); m.p.: 225-226 °C; ¹H(400 MHz, DMSO) 1.68-1.69 (m, 4H), 3.59 (t, *J*= 4.8 Hz, 2H), 4.08 (t, *J*= 6.4 Hz, 2H), 7.02 (s, 1H), 7.16 (s, 1H), 7.84 (dd, *J*= 8.0, 7.2 Hz, 2H), 8.15 (s, 1H), 8.42 (dd, *J*= 8.4, 0.8 Hz, 2H), 8.46 (dd, *J*= 7.2, 1.2 Hz, 2H), 12.4 (br s, 1H) ppm.
- [9] R.J. Neitz, S. Chen, F. Supek, V. Yeh, D. Kellar, J. Gut, et al. Lead Identification to Clinical Candidate Selection: Drugs for Chagas Disease. *J. Biomol. Screen.*, **2014**, 20: 101–111.
- [10] T. Schlecker, A. Schmidt, N. Dirdjaja, F. Voncken, C. Clayton, R.L. Krauth-Siegel. Substrate specificity, localization, and essential role of the glutathione peroxidase-type

- tryparedoxin peroxidases in *Trypanosoma brucei*. *J. Biol. Chem.*, **2005**, 280: 14385–94.
- [11] D. Sereno, J.L. Lemesre. Axenically cultured amastigote forms as an in vitro model for investigation of antileishmanial agents. *Antimicrob. Agents Chemother.*, **1997**, 41: 972–6.
- [12] D. Sereno, M. Cavaleyra, K. Zemzoumi, S. Maquaire, A. Ouaiissi, J.L. Lemesre. Axenically grown amastigotes of *Leishmania infantum* used as an in vitro model to investigate the pentavalent antimony mode of action. *Antimicrob. Agents Chemother.*, **1998**, 42: 3097–102.
- [13] A. Monks, D. Scudiero, P. Skehan, R. Shoemaker, K. Paull, D. Vistica, et al. Feasibility of a high-flux anticancer drug screen using a diverse panel of cultured human tumor cell lines. *J. Natl. Cancer Inst.*, **1991**, 83: 757–66.
- [14] J. Tavares, A. Ouaiissi, P. Kong Thoo-Lin, I. Loureiro, S. Kaur, N. Roy, et al. Bisnaphthalimidopropyl derivatives as inhibitors of *Leishmania* SIR2 related protein 1. *Chem Med Chem.*, **2010**, 5: 140–7.