



International Journal of Chemistry and Pharmaceutical Sciences

Journal Home Page: www.pharmaresearchlibrary.com/ijcps



Research Article

Open Access

Comparative Study for the Analysis of Cefixime trihydrate and its Degraded Products by Two RP-HPLC methods, one its Official and the other Developed Validated Method

Elsadig H. Kh. Adam^{*1} and Abdalfatah M. Bashir²

^{1,2}Department of Pharmaceutical Chemistry, College of Pharmacy, Prince Sattam bin Abdulaziz University, Riyadh, Al-kharj 11942, Saudi Arabia

ABSTRACT

The aim of the present work was aimed to carry out comparative study between method (1) [official method (BP, 2017)] and method (2) [developed validated analytical method (Adam *et al.*, 2012)] for the separation of cefixime trihydrate and its degraded products by using two different mobile phases, keeping the other parameters such as stationary phase, column condition, wavelength, and device constant. Mobile phase for method (1) consist of a solution of 0.03 M Tetra butyl ammonium hydroxide (pH 6.5) and acetonitrile with a ratio of 3:1 respectively while Mobile phase for method (2) consist of a mixture of 0.1M sodium dihydrogen phosphate monohydrate solution (pH 2.5) and methanol with a ratio 3:1 respectively. To study the degraded products sample was subjected to Sun light, UV light, and thermal effects. From data obtained proved the method (2) gave less retention time for the separation of drug with a larger number of decomposed products being detected compared by method (1).

Keywords: Cefixime trihydrate, Comparative Study, RP-HPLC method, decomposition of cefixime trihydrate, stability indicating method.

ARTICLE INFO

CONTENTS

1. Introduction	104
2. Materials and Method.	104
3. Results and Discussion.	104
4. Conclusion.	106
5. References	106

Article History: Received 18 February 2017, Accepted 25 March 2017, Available Online 27 April 2017

*Corresponding Author

Elsadig H. Kh. Adam
Department of pharmaceutical Chemistry,
College of Pharmacy,
Prince Sattam bin Abdulaziz University,
Riyadh, Al-kharj 11942, Saudi Arabia
Manuscript ID: IJCPS3344



PAPER-QR CODE

Citation: Elsadig H. Kh. Adam. Comparative Study for the Analysis of Cefixime trihydrate and its Degraded Products by Two RP-HPLC methods, one its Official and the other Developed Validated Method. *Int. J. Chem, Pharm, Sci.*, 2017, 5(4): 103-106.

Copyright © 2017 Elsadig H. Kh. Adam. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original work is properly cited.

1. Introduction

Cefixime is used to treat a wide variety of bacterial infections. This medication is known as a cephalosporin antibiotic. It works by stopping the growth of bacteria. Cefixime is a broad spectrum cephalosporin antibiotic and is commonly used to treat bacterial infections of the ear, urinary tract, and upper respiratory tract. The bactericidal action of Cefixime is because of the inhibition of cell wall synthesis. It binds to one of the penicillin binding proteins (PBPs) which inhibits the final transpeptidation step of the peptidoglycan synthesis in the bacterial cell wall, thus inhibiting biosynthesis and arresting cell wall assembly resulting in bacterial cell death. Cefixime is an orally active 3rd. generation cephalosporin which exerts its bactericidal action against both gram positive & gram negative organism by in bacterial cell wall synthesis. Chemically, Cefixime trihydrate name is (6R,7R)-7-[[2-(2-amino-1,3-thiazol-4-yl)-2-(carboxymethoxyimino)acetyl]amino]-3-ethenyl-8-oxo-5-thia-1-azabicyclo [4.2.0]oct-2-ene-2-carboxylic acid trihydrate (Figure1) and its molecular formula is $C_{16}H_{15}N_5O_7S_2 \cdot 3H_2O$ and molecular weight is 507.50 g/mol. It is a white powder that is freely soluble in water (1g/5ml) and stable in air, heat and acid solutions, while it is unstable in alkaline medium and light.




Figure 1: Structural formula of Cefixime Trihydrate

2. Materials and Methods

2.1 Materials (Chemicals and Reagents): Cefixime trihydrate was donated from DSM Company, all chemicals and reagents used were of a HPLC grade. Tetra butyl ammonium hydroxide 40% aqueous solution, Sodium dihydrogen phosphate monohydrate was obtained from AppliChem, Germany. Methanol and acetonitrile were obtained from fisher scientific UK Limited, UK. Water (HPLC gradient grade) supplied from Panreac, E.U. Orthophosphoric acid 85 % was obtained from BDH, England.

2.2 Instrument and Equipment:

- HPLC instruments a water Breeze 2 system, consisting of binary pump series 1525, UV/VIS detector 2489, and auto sampler series 2707.
- Sensitive balance, A&D Company limited, Japan.
- 827 pH lab. metrohm ion analysis, Herisau/ Switzerland.

2.3 Experimental:

2.3.1 Preparation of mobile phase for method (1) of 0.03 M Tetra butyl ammonium hydroxide solution (pH 6.5): Solution was prepared by weighing 8.2 g Tetra Butyl Ammonium hydroxide (or 20 ml of Tetra Butyl Ammonium hydroxide 40 % aqueous solution) and dissolving into 800

ml of distilled water and adjusted to pH 6.5 with %10 orthophosphoric acid and diluted up to 1000 ml with deionized water, and mixed with acetonitrile with a ratio of 4:1 respectively and degassed (BP., 2017).

2.3.2 Preparation of mobile phase for method (2) of 0.1M sodium dihydrogen phosphate monohydrate solution (pH 2.5): Solution was prepared by weighing 13.67 g of sodium dihydrogen phosphate monohydrate and dissolving into 900 ml of deionized water and adjusted to pH 2.5 with diluted orthophosphoric acid and diluted up to 1000 ml with distilled water, and mixed with methanol with a ratio of 3:1 respectively and degassed (Adam *et al.*, 2012).

2.3.3 Chromatographic conditions used for the analysis of cefixime trihydrate and its degraded products:

- Mobile phase: (1) & (2)
- Flow Rate: 1.0 ml/min
- Injection volume = 50 μ l
- Column = Waters Spherisorb@5.0 μ m ODS2 250 mm x4.6 mm ID.
- Temperature = room temperature (Ambient)
- Detection wave length at 254 nm

2.3.4 Preparation of standard stock solution: Stock standard solution having concentration 100 μ g/ml was prepared by dissolving pure drug of cefixime trihydrate in water, injected into the chromatographic column, (figure (2) & (3) and table 1.

2.3.5 Preparation of degradation products of cefixime:

2.3.5.1 Preparation of the decomposed product cefixime trihydrate solid by sun-light

About 5.0 grams of cefixime trihydrate solid were placed between two glass plates (20 x 20 cm), sealed with gum tape and directly exposed to sunlight for six months(March to August). Samples were taken every month and tested for degradation by HPLC (table .2).

2.3.5.2 Preparation of the decomposed product cefixime trihydrate solution by UV-light

100 μ g / ml of cefixime trihydrate solution in water were prepared and transferred to a stoppered tube. The solutions were placed under UV radiation at 254 nm. Samples were taken at 30, 60, 90, 120, and 150 minutes and tested for degradation by HPLC (table .3).

2.3.5.3 Preparation of the decomposed product cefixime trihydrate solid by thermal at 100 °C

Five grams of CEF-3H₂O solid were placed in a petri dish and was put in oven at 100 °C. Samples were taken every hour and tested for degradation by HPLC (Table.4).

2.3.5.4 Preparation of the decomposed cefixime trihydrate solution by thermal at 100 °C for 45 minutes.

Solution of cefixime trihydrate (10 mg / 100 ml water) was prepared. The flask was placed into a water-path thermostatic at 100 °C for 45 minute table 5.

3. Results and Discussion

Under optimization condition for the RP-HPLC method (1) and method (2) with keeping others fixed and changeable mobile phase ,the separation chromatogram obtained of cefixime trihydrate reference standard was appear in figure

(2 & 3), it was found that cefixime trihydrate separation at 9.75 minute by using method (2) while separation at 13.03 minute by using method (1). Discrimination developed validated method (2) (Adam *et al.*, 2012) versus method (1) (BP., 2017), illustrated that from the assay test, the retention time of CEF-3H₂O less than that time obtained by method (1) were 9.75, 13.03 minutes respectively, (Table 1) and figure 4. There's very little noticeable change when testing for degradation of cefixime trihydrate solid under the influence of sunlight by using the analysis methods (1) & (2) table 2 and figure 5. From the results obtained for the tested degraded products under stress condition of UV for cefixime trihydrate solution, two methods were given two degradation products but better resolution by the method (2), table 3 and figure 6 & 7. The analysis of cefixime trihydrate solid thermal decomposed at 100 °C by using method (1) & (2), revealed that method (2) detected three decomposed products while method (1) was detect only two decomposed, Table 4, and Figures 8&9. There is no noticeable quantitatively and qualitatively change for the analysis of cefixime trihydrate solution thermal decomposed by using method (1) & (2).




Figure 2




Figure 3

Table 1: Analysis of cefixime trihydrate reference standard by method (1) & (2).

Parameters	Retention Time R.T/minutes
Method 1	13:01
Method 2	9.75

Table 2: Analysis of decomposed cefixime trihydrate solid form by sun- light using method (1) & (2).

Interval time/month	Method (1) % Remaining content	Method (2) % Remaining content
1 Month	79.72	78.56
2 Months	64.71	64.51
3 Months	47.45	48.05
4 Months	22.99	22.57
5 Months	11.62	11.65
6 Months	8.11	8.15

Table 3: Analysis of decomposed cefixime trihydrate solution by UV-light using method (1) & (2).

Results	method (1)		method (2)	
	Content %	R.T	Content %	R.T
Decomposed (1)	17.46%	4.62	18.83%	2.97
Decomposed (2)	23.34%	11.05	22.88%	15.04
Remaining CEF-3H ₂ O	26.89%	13.03	27.19%	9.75

Table 4: Analysis of decomposed cefixime trihydrate solid by thermal effect at 100 °C using method (1) & (2).

Results	Method (1)		Method (2)	
	Content %	R.T	Content %	R.T
Decomposed (1)	8.50%	4.51 min	7.89%	3.91 min
Decomposed (2)	12.71%	6.38 min	14.86%	4.216 min
Decomposed (3)	-	-	8.20%	5.899 min
Remaining CEF-3H ₂ O	71.06%	13.014 min	66.34%	9.750 min

Table 5: Analysis of decomposed cefixime trihydrate solution by thermal effect at 100°C using method (1) & (2).

Results	method (1)		method (2)	
	Content %	R.T	Content %	R.T
Decomposed (1)	8.90%	1.78 min	7.86%	14.07 min
Decomposed (2)	37.61%	11.05 min	36.97%	14.99 min
Remaining CEF-3H ₂ O	51.43%	13.02 min	51.69%	9.70 min

4. Conclusion

This work described the evaluation of analytical method (1) & (2). The method (2) described enables the quantification and qualification of cefixime trihydrate and its degraded products compared to the method (1). The data obtained demonstrate good precision proves the reliability of the method (2) and separation of the degradation products did not interfere with the active ingredient cefixime trihydrate. Hence, the method (2) can be used routinely for qualitative and quantitative estimation of cefixime trihydrate and it can also be used as stability indicating method.

5. References

- [1] Adam, Elsadig H. K; Saeed, Ahmed E. M; & Barakat, Izzeldin E. Development and Validation of a High Performance Liquid Chromatography Method for Determination of Cefixime Trihydrate, IJPSR. International journal of Pharmaceutical sciences and research, 2012; Vol. 3(2): 469-473.
- [2] Amin MI and Bryan JT, (1973) .Kinetic factors affecting stability in aqueous formulation. *pharm. Sc.* 62 (11):1768-1771.
- [3] Analytical chemistry, Six Edition, Gary D. Christian, John Wiley & Sons, Inc., 2003, USA.
- [4] British Pharmacopoeia Commission (2017) the British Pharmacopoeia, Volume I, the Stationary Office Ltd., London, England, p. 451.
- [5] Brogden, R.N.; Richards, D.M. Cefixime: a review of its antibacterial activity, pharmacokinetic properties and therapeutic potential. *Drugs*, v.38, n.4, p.524-550, 1989.
- [6] Dhoka, M.V.; Sandage, S.J.; Dumbre, S.C. Simultaneous determination of cefixime trihydrate and dicloxacillin sodium in pharmaceutical dosage form by reversed-phase high-performance liquid chromatography. *J. AOAC Int.*, v.93, n.2, p.531-535, 2010.
- [7] Food and drug Administration (FDA, 2000) "Analytical Procedures and Methods Validation: Chemistry, Manufacturing and Controls Documentation; Availability," Federal Register (Notices) 65(169), 52776–52777.
- [8] Gonzalez-Hernandez, R.; Nuevas-Paz, L.; Soto-Mulet, L.; Lopez-Lopez, M.; Hoogmartens, J. Reversed phase high performance liquid Chromatographic determination of cefixime in bulk drugs. *J. Liq. Chromatogr. Relat. Technol.*, v.24, n.4, p.2315-2324, 2001.
- [9] International Conference on Harmonization. (ICH). International Conference on Harmonization of technical requirements for registration of pharmaceuticals for human use. Quality Guidelines. Q2 Analytical Validation. Q2 (R1). Validation of analytical procedures: text and methodology. ICH, 2005.
- [10] International Conference on Harmonization (ICH, 2005). Tripartite Guideline
- [11] Validation of analytical procedure text and methodology, Q2 [R1].
- [12] Khandagle, K.S.; Ganddhi, S.V.; Deshpande, P.B.; Gaikwad, N.V. A simple and sensitive RPHPLC method for simultaneous estimation of cefixime and ofloxacin in combined tablet dosage form. *Int. J. Pharm. Pharm. Sci.*, v.3, n.1, p.46-48, 2011.
- [13] Martindale (2009), Thirty –Sixth edition, Edited by Sean C Sweetman. Royal Pharmaceutical Society of Great Britain (RPS) Publishing, UK.
- [14] Meyer .V.R, (1994). Practical high performance liquid chromatography. J.Wiley and Sons .Chichester. London.p 10-25.
- [15] Shabir G.A., (2003), "Validation of HPLC Chromatography Methods for Pharmaceutical Analysis. Understanding the Differences and Similarities between validation requirements of FDA, the US Pharmacopeia and the ICH," *J.Chromatogr. A.* 987(1-2), 57-66.
- [16] The United States pharmacopeia 33-National formulary 28, Vol I, validation of Compendial procedures (1225) .unit state pharmacopeia Convention on (2010) P 774-77.
- [17] The united States Pharmacopeia 34- National Formulary 29, (2011). Vol II, united State pharmacopeia convention on (2011); p 2220.
- [18] Zendelovska D., Stafilov T., Melosevski P, (2003). High performance liquid Chromatographic method for determination of cefixime and cefotxime in human plasma *Bull. Chem. Tech. of Macedoni*, p.2221-39.