



# International Journal of Medicine and Pharmaceutical Research

CODEN (USA): IJCPNH | ISSN: 2321-2624  
Journal Home Page: [www.pharmaresearchlibrary.com/ijmpr](http://www.pharmaresearchlibrary.com/ijmpr)



## In-Vitro Cytotoxic Activity of *Persea Americana* Miller Leaves Extracts

Narayan R Miskin\*<sup>1</sup>, Manjunatha S Katagi<sup>1</sup>, Girish Bolakatti<sup>2</sup>, Vishvakarma kamala Sunita<sup>1</sup>, Suchitra.M<sup>1</sup>

<sup>1</sup>Bapuji Pharmacy College, SS Layout, Davangere-577004, Karnataka, India.

<sup>2</sup>GM Institute of Pharmaceutical Sciences and Research, Davangere-577006, Karnataka, India.

### ABSTRACT

Phytochemistry plays an important role for curing of various human diseases with slight side effects. Such medicinal plants have bioactive compounds may be primary or secondary metabolites. The present study aims to evaluate that the In-vitro anticancer activity of *Persea americana* miller leaves extracts in different cancer cell lines. Dried Coarse powered leaves of *Persea americana* Miller was successively extracted with various solvents based on their polarity such as petroleum ether, chloroform, ethyl acetate and aqueous extract by maceration. Qualitative phytochemical screening was performed to detect phytochemical constituents of the extracts. Cytotoxic activity of plant leaves extracts on MCF-7 and HeLa cells was investigated *in vitro* through MTT bioassay. All the extracts except EAPA has showed IC<sub>50</sub> value of 233.7µg/ml, 182.0µg/ml, 203.7µg/ml, 147.7µg/ml, 121.0µg/ml respectively in MCF-7 cells. All the extracts have showed IC<sub>50</sub> value 200.2µg/ml, 105.4µg/ml, 285.8µg/ml, 132.7µg/ml and 214.4µg/ml respectively in HeLa cells.

**Keywords:** *Persea americana* miller, cytotoxic activity, MTT assay.

### ARTICLE INFO

#### \*Corresponding Author

Narayan R Miskin  
Bapuji Pharmacy College,  
SS Layout, Davangere-577004,  
Karnataka, India.



**ARTICLE HISTORY:** Received 25 Dec 2019, Accepted 29 Feb 2020, Available Online 10 April 2020

©2020 Production and hosting by International Journal of Medicine and Pharmaceutical Research. All rights reserved.

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original work is properly cited.

**Citation:** Narayan R Miskin, et al. Structurally In-Vitro Cytotoxic Activity of *Persea Americana* Miller Leaves Extracts. *Int. J. Med. Pharm. Res.*, 2020, 8(2): 58-62.

### CONTENTS

1. Introduction. . . . .	58
2. Materials and Methods. . . . .	59
3. Results and Discussion. . . . .	59
4. Acknowledgement. . . . .	61
5. References . . . . .	61

### 1. Introduction

Cancer is characterized by uncontrolled division of cells and the ability of these cells to invade other tissues either by direct growth into adjacent tissues through invasion or by implantation into distant sites by metastasis<sup>1</sup>. Cancer is the leading cause of death in economically developed countries and the second leading cause of death in

developing countries and the burden of cancer is increasing in economically developing countries as a result of<sup>2</sup> population aging and growth as well as increasingly an adoption of cancer-associated lifestyle choices including smoking, physical inactivity, and “westernized” diets<sup>3</sup>. It has been estimated that the total number of new cases of cancer will rise from 10 million in year 2000 by

approximately 25% in each decade, reaching 24 million new cases per year in the year 2050; the total number of deaths will rise from 6 million in the year 2000 to 10 million in 2020 to over 16 million in the year 2050; in the year 2050, there will be 17 million new cases of cancer in less developed countries, while only 7 million new cases of cancer will occur in the more developed countries<sup>4,5,6</sup>.

In chemotherapeutic cancer natural or synthetic biological or chemical agents are used to reverse, suppress or prevent carcinogenic progression<sup>5</sup>. Historically natural products have been employed as anticancer agents for quite a considerable period of human existence and through the years have been incorporated into both traditional and allopathic medicine. A significant number of chemotherapeutic drugs in current use were either isolated from plant species or derived from natural prototype<sup>2,6</sup>. The following are some of excellent examples of vinca alkaloids, vincristine, vinblastine isolated from catharanthus roseus, etoposides and teneposide, the semi synthetic derivative of epipodophylotoxin, isolated from *Podophyllum* species and the naturally derived taxanes isolated from taxus species and semisynthetic derivative of camptothecin, irinotecan and topotecan, isolated from *Camptotheca accuminata*<sup>7,8</sup>. In fact, according to Cragg and Newman, more than 50% of drugs in clinical trials from natural sources or are related to them<sup>9</sup>.

Plant materials have a long history of used in the treatment of cancer. Hartwell has reported in his review about the 300 plants species used against<sup>10, 11, 13</sup>. Plant based discovery has resulted in the development of many anticancer drugs currently in clinical use. Besides this it also provides a platform for the design of novel and safe drugs through a proper understanding of the complete synergistic interaction of various constituents of anticancer herbs<sup>14, 15, 16</sup>

*Persea americana* (family Lauraceae) is a tree plant known as 'avocado', 'avocado pear' or 'alligator pear'. They are widely cultivated throughout the tropics and subtropics of the world for their edible fruits and for some economic and therapeutic uses<sup>17</sup>. Avocado tree is known only for the fruit that people usually consume. Apparently avocado leaf is one of the natural ingredients that can be used as a traditional medicine<sup>18</sup>. This leaf has been empirically used as a diuretic, analgesic, anti-inflammatory, hypertensive, hypoglycemic, diarrhea, sore throat and hemorrhage cure.

Avocado is one of the fruit plant groups which are nutritious as a preservative and antioxidants<sup>19,20</sup>. Avocado flesh can be used as an anti-hyperlipidemia and has antioxidant activity and reduce the risk of metabolic syndrome<sup>21,22,23</sup>. One part of the avocado plant that has the potential as a natural antioxidant substance is avocado leaf. Previous research has shown that avocado leaf has the potential as a natural antioxidant<sup>24</sup> and positively contains alkaloids, flavonoids, saponins, tannins and steroids using methanol solution to hydrolyze and extract avocado leaf<sup>25</sup>. The present study was aimed to investigate the *in vitro* cytotoxic potential of Petroleum ether, chloroform,

ethyl acetate and aqueous extracts of leaves of *Persea americana* miller against MCF-7 and HeLa cancer cells by MTT bioassay.

## 2. Material and Methods

### Collection of plant material and extraction:

The leaves of plant *Persea americana* Miller used for the present study were collected from the local surroundings of Davangere and authenticated by professor L.C. Kulkarni, Department of botany, P.C.Jabin science college, Hubballi. Authenticated leaves of *Persea americana* Miller. were shade dried and pulverized into coarse material. #120gms per batch coarsely dried powdered leaves of *Persea americana* Miller were successively extracted with various solvents in increasing order of polarity such as petroleum ether, chloroform, ethyl acetate and macerating with water. Percentage yield of various extracts of *Persea americana* Miller leaves were determined. Preliminary phytochemical analysis is carried out to assess the presence of various phytoconstituents present in plant leaves extracts.

### *In vitro* anticancer activity by MTT Bioassay

#### Cytotoxicity studies for MCF-7 and HeLa cell lines

##### Cell lines and culture medium:

All the cell lines were procured from ATCC stock cells was cultured in DMEM/RPMI. The monolayer cell culture was trypsinized and the cell count was adjusted to  $5.0 \times 10^5$  cells/ml using respective media containing 10% FBS. To each well of the 96 well microtiter plate, 100  $\mu$ l of the diluted cell suspension (50,000cells/well) was added. After 24 h, when a partial monolayer was formed, the supernatant was flicked off, washed the monolayer once with medium and 100  $\mu$ l of different test concentrations of test drugs were added on to the partial monolayer in microtiter plates. The plates were then incubated at 37°C for 24hrs in 5% CO<sub>2</sub> atmosphere. After incubation the test solutions in the wells were discarded and 100  $\mu$ l of MTT (5 mg/10 ml of MTT in PBS) was added to each well. The plates were incubated for 4 h at 37° C in 5% CO<sub>2</sub> atmosphere. The supernatant was removed and 100  $\mu$ l of DMSO was added and the plates were gently shaken to solubilize the formed formazan. The absorbance was measured using a microplate reader at a wavelength of 590 nm. The percentage growth inhibition was calculated using the following formula and concentration of test drug needed to inhibit cell growth by 50% (IC<sub>50</sub>) values is generated from the dose-response curves for each cell line<sup>30,31</sup>.

##### Calculating Inhibition:

$$\% \text{ Inhibition} = ((\text{OD of Control} - \text{OD of sample}) / \text{OD of Control}) \times 100.$$

## 3. Results and Discussion

Successive extraction with different solvents and their % yield was reported in Table no 1. phytochemical screening of all extracts were done and reported their phytochemical constituents in Table no2. Cytotoxicity activity of four extracts were carried out against MCF 7 cell line and HeLa cell line at different concentrations to determine the IC<sub>50</sub> (50% growth inhibition) by MTT assay. Results of different concentrations of *Persea americana* Miller including 10 – 320ug/ml are tabulated in Table no 3, and graphically





